

柑橘類蟲害防治藥劑一覽表

| 農藥名稱 | 稀釋倍數 或施藥量 | 安全 採收期 | 有殼介殼蟲 | 無殼介殼蟲 | 粉介殼蟲 | 蚜蟲類 | 木蟲、刺粉蟲 | 潛葉蛾 | 葉蟎 | 銹蟎 | 椿象類 | 星天牛 | 小黃薊馬 |
|------------------------|--------------|-----------|-------|-------|------|-----|--------|-----|----|----|-----|-----|------|
| 納得護賽寧 33%可濕性粉劑 | 1,200 | 9 天 | ● | | | | | | | | | | |
| 馬拉松 50%乳劑 | 800 | 14 天 | ● | ● | ● | ● | ● | ● | | | | | |
| 馬拉松 50%乳劑 +夏油 95%乳劑 | 800+95 | 14 天 | ● | ● | ● | ● | ● | ● | ● | ● | | | |
| 大滅松 20%乳劑 | 450 | 21 天 | ● | ● | ● | ● | ● | ● | | | ● | | |
| 大滅松 44%乳劑 | 1,000 | 21 天 | ● | ● | ● | ● | ● | ● | | | ● | | |
| 布芬三亞蟎 20%乳劑 | 1,000 | 18 天 | ● | | | | | | | | | | |
| 丁基加保扶 31.6%乳劑 | 400 | 21 天 | ● | | | ● | | | | | | | |
| 夏油 95%乳劑 | 95 | 免訂 | ● | ● | | | | | ● | | | | |
| 礦物油 99%乳劑 | 500 | 免訂 | | | | | | | ● | | | | |
| 滅大松 40%乳劑 | 1,000 | 18 天 | | ● | | | | | | | | | |
| 加福松 50%乳劑 | 1,200 | 15 天 | | | ● | | | | | | | | |
| 加保扶 44%水懸劑 | 1,200 | 7 天 | | | | ● | ● | | | | | | |
| 加保扶 40.64%水懸劑 | 1,200 | 7 天 | | | | ● | ● | | | | | | |
| 滅賜松 25%乳劑 | 1,000 | 14 天 | | | | ● | | ● | | | | | |
| 硫滅松 25%乳劑 | 1,000 | 18 天 | | | | ● | | | | | | | |
| 免扶克 20%乳劑 | 800 | 20 天 | | | | ● | | | | | | | |
| 丁基加保扶 31.6%乳劑 | 1,200 | 20 天 | | | | ● | | | | | | | |
| 丁基加保扶 40%可濕性粉劑 | 1,200 | 20 天 | | | | ● | | | | | | | |
| 丁基加保扶 48.34%乳劑 | 1,000 | 21 天 | | | | | | | | | | | ● |
| 陶滅蝨 50%可濕性粉劑 | 1,000 | 21 天 | | | | | ● | | | | | | |
| 納得亞滅寧 13.5%乳劑 | 1,000 | 6 天 | | | | | | ● | | | | | |
| 陶斯松 50%可濕性粉劑 | 1,500 | 7 天 | | | | | | ● | | | | | |
| 護賽寧 5%溶液 | 1,500 | 9 天 | | | | | | ● | | | | | |
| 益滅賽寧 40%可濕性粉劑 | 2,500 | 12 天 | | | | | | ● | | | | | |
| 芬諾克 25%可濕性粉劑 | 2,000 | 12 天 | | | | | | ● | | | | | |
| 陶斯寧 25%乳劑 | 1,000 | 14 天 | | | | | | ● | | | | | |
| 賽洛寧 1%可濕性粉劑 | 1,600 | 14 天 | | | | | | ● | ● | | | | |
| 賽洛寧 2.8%乳劑 | 4,000 | 14 天 | | | | | | ● | | | | | |
| 賽洛寧 2.5%微乳劑 | 4000 | 14 天 | | | | | | ● | | | | | |
| 賽洛寧 2.46%膠囊懸著液 | 4000 | 14 天 | | | | | | ● | | | | | |

| 農藥名稱 | 稀釋倍數 或施藥量 | 安全採 收期 | 有殼介殼蟲 | 無殼介殼蟲 | 粉介殼蟲 | 蚜蟲類 | 木蟲、刺粉蟲 | 潛葉蛾 | 葉蟎 | 銹蟎 | 椿象類 | 星天牛 | 小黃薊馬 |
|---------------|--------------|-----------|-------|-------|------|-----|--------|-----|----|----|-----|-----|------|
| 培丹 50%可溶性粉劑 | 1,000 | 14 天 | | | | | | ● | | | | | |
| 硫滅松 25%乳劑 | 500 | 18 天 | | | | | | ● | | | | | |
| 益滅松 50%可濕性粉劑 | 500 | 18 天 | | | | | | ● | | | | | |
| 免扶克 40%水懸劑 | 1,200 | 20 天 | | | | | | ● | | | | | |
| 畢芬寧 2.8%乳劑 | 1,000 | 20 天 | | | | | | ● | | | | | |
| 畢芬寧 2.5%水懸劑 | 1,000 | 20 天 | | | | | | ● | | | | | |
| 納乃得 40%可溶性粉劑 | 1,500 | 20 天 | | | | | | ● | | | | | |
| 納乃得 40%水溶性粒劑 | 1,500 | 20 天 | | | | | | ● | | | | | |
| 納乃得 25%可溶性粒劑 | 900 | 20 天 | | | | | | ● | | | | | |
| 納乃得 24%溶液 | 750 | 20 天 | | | | | | ● | | | | | |
| 福隆納乃得 30%水懸劑 | 1,500 | 20 天 | | | | | | ● | | | | | |
| 佈飛賽滅寧 25%乳劑 | 1,000 | 21 天 | | | | | | ● | | | | | |
| 毆殺滅 10%溶液 | 350 | 7 天 | | | | | | ● | | | | | |
| 密滅汀 1%乳劑 | 1,500 | 6 天 | | | | | | | ● | | | | |
| 依殺蟎 10%水懸劑 | 4,000 | 6 天 | | | | | | | ● | | | | |
| 畢達本 20%可濕性粉劑 | 3,000 | 6 天 | | | | | | | ● | ● | | | |
| 合賽多 5%乳劑 | 2,000 | 9 天 | | | | | | | ● | | | | |
| 芬普寧 10%乳劑 | 1,500 | 10 天 | | | | | | | ● | | | | |
| 畢芬寧 2.5%水懸劑 | 750 | 10 天 | | | | | | | ● | | | | |
| 畢芬寧 2.8%乳劑 | 750 | 10 天 | | | | | | | ● | | | | |
| 扶吉胺 39.5%水懸劑 | 2,000 | 10 天 | | | | | | | ● | | | | |
| 芬普毆蟎多 13%水基乳劑 | 1,000 | 15 天 | | | | | | | ● | | | | |
| 芬普蟎 5%水懸劑 | 2,000 | 15 天 | | | | | | | ● | | | | |
| 佈賜芬蟎 45%水懸劑 | 2,500 | 15 天 | | | | | | | ● | | | | |
| 達馬芬普寧 45%乳劑 | 1,500 | 15 天 | | | | | | | ● | | | | |
| 合賽多 10%可濕性粉劑 | 4,000 | 20 天 | | | | | | | ● | | | | |
| 芬硫克 35%乳劑 | 1,500 | 20 天 | | | | | | | ● | | | | |
| 芬殺蟎 10%乳劑 | 1,600 | 21 天 | | | | | | | ● | | | | |
| 合賽芬普寧 7.5%乳劑 | 1,500 | 21 天 | | | | | | | ● | | | | |
| 愛殺松 46.5%乳劑 | 800 | 21 天 | | | | | | | ● | | | | |
| 畢汰芬 4%水懸劑 | 3,000 | 25 天 | | | | | | | ● | | | | |
| 新殺蟎 25%乳劑 | 500 | 30 天 | | | | | | | ● | | | | |

| 農藥名稱 | 稀釋倍數 或施藥量 | 安全 採收期 | 有殼介殼蟲 | 無殼介殼蟲 | 粉介殼蟲 | 蚜蟲類 | 木蝨、刺粉蝨 | 潛葉蛾 | 葉蟎 | 銹蟎 | 椿象類 | 星天牛 | 小黃薊馬 |
|-----------------|--------------|-----------|-------|-------|------|-----|--------|-----|----|----|-----|-----|------|
| 蟎離丹 25%可濕性粉劑 | 500 | 30 天 | | | | | | | ● | | | | |
| 三亞蟎 20%乳劑 | 1,000 | 30 天 | | | | | | | ● | | | | |
| 大克蟎 18.5%乳劑 | 400 | 30 天 | | | | | | | ● | | | | |
| 大克蟎 18.5%可濕性粉劑 | 400 | 30 天 | | | | | | | ● | | | | |
| 大克蟎 35%乳劑 | 800 | 30 天 | | | | | | | ● | | | | |
| 大克蟎 35%可濕性粉劑 | 750 | 30 天 | | | | | | | ● | | | | |
| 大克蟎 42%乳劑 | 900 | 30 天 | | | | | | | ● | | | | |
| 芬佈賜 50%可濕性粉劑 | 2,000 | 30 天 | | | | | | | ● | | | | |
| 毆蟎多 30%可濕性粉劑 | 1,000 | 30 天 | | | | | | | ● | | | | |
| 毆蟎多 57%乳劑 | 1,500 | 30 天 | | | | | | | ● | | | | |
| 克芬蟎 42%水懸劑 | 3,000 | 30 天 | | | | | | | ● | ● | | | |
| 得芬瑞 10%可濕性粉劑 | 3,000 | 9 天 | | | | | | | ● | | | | |
| 賜派芬 30%水懸劑 | 5,000 | 6 天 | | | | | | | ● | | | | |
| 可濕性硫磺 95%可濕性粉劑 | 100 | | | | | | | | | ● | | | |
| 可濕性硫磺 80%水分散性粒劑 | 85 | | | | | | | | | ● | | | |
| 可濕性硫磺 90%可濕性粉劑 | 96 | | | | | | | | | ● | | | |
| 可濕性硫磺 75%水分散性粒劑 | 80 | | | | | | | | | ● | | | |
| 阿巴汀 2%乳劑 | 2000 | 6 天 | | | | | | | | ● | | | |
| 毆殺滅 10%溶液 | 350 | 7 天 | | | | | | | | ● | | | |
| 丁基加保扶 25%可濕性粉劑 | 1,000 | 12 天 | | | | | | | | ● | | | |
| 汰芬隆 50%可濕性粉劑 | 1,000 | 15 天 | | | | | | | | ● | | | |
| 加福松 50%乳劑 | 1,500 | 15 天 | | | | | | | | ● | | | |
| 福隆納乃得 30%水懸劑 | 1,000 | 20 天 | | | | | | | | ● | | | |
| 芬普蟎 5%水懸劑 | 1,000 | 21 天 | | | | | | | | ● | | | |
| 新殺蟎 25%乳劑 | 1,000 | 21 天 | | | | | | | | ● | | | |
| 新殺福化利 50%乳劑 | 2,000 | 21 天 | | | | | | | | ● | | | |
| 芬佈賜 50%水懸劑 | 2,000 | 60 天 | | | | | | | | ● | | | |

| 農 藥 名 稱 | 稀釋倍數 或施藥量 | 安全 採收期 | 有 殼 介 殼 蟲 | 無 殼 介 殼 蟲 | 粉 介 殼 蟲 | 蚜 蟲 類 | 木 蝨 、 刺 粉 蝨 | 潛 葉 蛾 | 葉 蟎 | 銹 蟎 | 椿 象 類 | 星 天 牛 | 小 黃 薊 馬 | 東 方 果 實 蠅 |
|----------------|--------------|-----------|-----------------------|-----------------------|------------------|-------------|----------------------------|-------------|--------|--------|-------------|-------------|------------------|-----------------------|
| 加保利 50%可濕性粉劑 | 500 | 18 天 | | | | | | | | | ● | | | |
| 加保利 85%可濕性粉劑 | 850 | 18 天 | | | | | | | | | ● | | | |
| 大滅松 20%乳劑 | 450 | 21 天 | | | | | | | | | ● | | | |
| ◎加保扶 40.64%水懸劑 | 100 | 10 天 | | | | | | | | | | ● | | |
| ◎加保扶 44%水懸劑 | 100 | 10 天 | | | | | | | | | | ● | | |
| 芬化利 20%乳劑 | 2,000 | 9 天 | | | | | | | | | | | | ● |
| 芬化利 20%水基乳劑 | 2,000 | 9 天 | | | | | | | | | | | | ● |
| 芬殺松 50%乳劑 | 1,000 | 21 天 | | | | | | | | | | | | ● |
| ◎賜諾殺 0.02%餌劑 | 8 | | | | | | | | | | | | | ● |

◎加保扶僅使用於樹幹基部噴灑防治星天牛。

◎賜諾殺餌劑僅可以誘殺器懸掛進行誘殺，不可噴及作物。

※表列植物保護手冊柑桔登記之防治藥劑如有更新，請自行上網查詢 (<http://www.tndais.gov.tw/Protect/plant.htm>，植物保護手冊(連結至農業藥物毒物試驗所))。