

貯藏期間稻米品質的改變¹

王聯輝 謝順景

國立中興大學、臺中區農業改良場

摘要

稻米之理化性隨貯藏條件及貯藏期間之不同，而發生不同程度之改變。本文就貯藏後米粒老化的一般機制，貯藏對米粒外貌及碾米品質的影響，貯藏對米粒的烹調品質的影響，貯藏對食用品質的影響，及貯藏對米粒營養品質的影響加以檢討。

游離脂肪酸隨時間之延長而增加，並形成脂肪酸——直鏈澱粉複合物、pH 降低，因而加速老化。膠化溫度隨貯藏期間之延長而升高，加熱吸水率及膨脹體積亦因溫度及貯藏時間的增加而增加。米飯之膠體軟硬度隨貯藏時間之延長而變硬，粘性則降低。貯藏時間愈長則游離胺基酸及維生素 B₁ 之含量降低愈多，但蛋白質含量之變化不多。

前言

近年來臺灣稻米在各有關單位的努力之下，糙米年產量一直維持在 240 萬公噸左右，加上國民消費型態的改變，稻米每年人消費量由 1980 年之 105.5 kg 降至 1982 年之 95 kg，致使每年過剩稻米高達 50 萬公噸以上⁽³⁾。因消費量有限，所以稻米的貯藏期大都延長了⁽¹⁾，在貯藏期間，由於受到穀粒胚部本身的呼吸作用以及昆蟲、微生物的為害，所釋放的熱及水氣，使得稻米的溫度及周圍環境的相對濕度升高⁽⁷⁾，而影響米粒的外貌 (grain appearance)^(2,1,17,21)、碾米品質 (milling quality)⁽¹⁷⁾，烹調品質 (cooking quality)^(16,17)、食用品質 (eating quality)^(16,10,11)，以及營養品質 (nutritional quality)^(8,9,20)。由於臺灣的產米過剩，稻穀倉儲期間較長，同時無冷藏設備，故經貯藏 2 ~ 3 年後品質變劣，即白米的透明度隨貯存期的延長而變差，黃變米率亦增加，白米之碱性擴散度 (alkali spreading) 膠體展流度 (gel consistency) 及食味品質亦降低⁽²⁾，如何改善貯藏之條件及維持稻米之品質為極需解決的問題。本文針對貯藏期間稻米品質的一些變化情形，作較深入的探討。

一、貯藏後米粒老化的一般機制

稻米米粒中成分的含量因品種及栽培環境而有所不同^(10,11)。一般在水分含量為 14% 時，米粒中澱粉約佔穀粒乾重的 70~80%，蛋白質約佔 6~9%，脂肪 (lipid) 約有 2~3%⁽⁸⁾ (表 1)。而這三種主要物質經貯藏後，由於發生水解或氧化作用，而影響米粒的外觀及烹調品質，進而更影響到米飯的香味 (aroma) 及質地 (Texture)。Moritaka and Yasumatsu⁽¹⁰⁾ 提出了米粒老化的機制 (表 2)，認為米粒貯藏後，脂肪經水解後形成之游離脂肪酸 (free fatty acid) 能與澱粉中之直

1. 臺中區農業改良場研究報告第 0166 號。

表一、糙米、白米和米糠之化學組成含量

Table 1. Chemical composition (dry basis) of brown rice, milled rice, and bran
 (Resurreccion *et al* 1979; Singh and Juliano 1977; Cagampang *et al*
 1976).

Constituent	Brown rice (0-100%) ^a	Milled rice (90-100%) ^a	Bran (0-6%) ^a
Starch (% anhydroglucose)	75.9	89.8	9.7
Amylose (%)	30.8	32.7	6.7
Total sugars (% glucose)	1.3	0.4	6.4
Crude fiber (%)	0.8	0.1	9.7
Dietary fiber (%)	1.9	0.7	28.4
Crude fat (%)	3.3	0.6	22.8
Crude protein (%)	8.4	7.7	15.7
Crude ash (%)	1.5	0.56	10.6
Phosphorus (%)	0.27	0.09	1.7
Phytin phosphorus (%)	0.14	0.03	1.1
Iron (mg/100 g)	2.0	0.67	15.7
Zinc (mg/100 g)	2.1	1.3	10.9
Lysine (g/16g N)	4.1	3.8	5.6
Threonine	3.7	3.7	4.1
Methionine+cystine (g/16g N)	4.7	4.9	4.7
Tryptophan (g/16g N)	1.2	1.2	1.2

^aAbrasive milling fraction of brown rice (% by wt)。


鏈性澱粉 (amylose) 結合成爲脂肪酸一直鏈性澱粉 fatty acid-amylose 的複合物，而抑制了澱粉顆粒的膨脹 (swelling)，而使米飯的質地變硬。另一方面，脂肪或是游離脂肪酸，經氧化後可產生氫過氧化物 (hydroperoxides) 及羰基化合物 (carbonyl compounds) 這二種物質可促進蛋白質的氧化，使得蛋白質中之-S-H 基相互作用產生了 S-S 基的雙硫鍵結，可影響米粒的外貌、降低揮發性硫化物的含量，另外也會抑制澱粉顆粒的膨脹，而影響米飯的香味及質地。關於澱粉的改變，可能是經過貯藏後，澱粉顆粒中之微束鍵結 (micelle binding) 加強了，而使米粒質地變硬。

二、貯藏對米粒外貌及碾米品質的影響

稻米採收以後，若溫度在15°C以上貯藏3~4個月後，其品質就開始發生變化⁽¹⁷⁾。其中米粒外貌的改變主要是白度 (whiteness) 及透明度 (translucency) 會隨貯藏溫度及期間的增加而降低 (表3、表4)；黃米率則隨之增加^(1,2,17) (圖1)。

表二、米粒老化的機制

Table 2. Summary of Mechanisms of Aging of Rice Grain*



* Source: Moritaka and Yasumatsu (1972b); used by permission.

表三、不同貯藏條件下 (2°C及29°C) 貯藏 6 個月後之白米其硬度和白色度之變化

Table 3. Changes in the hardness index and whiteness of milled rice stored in various forms at 2°C and 29°C for 6 months.

Type of rice and form stored	Crop season	Hardness index*		Whiteness ^b (%)	
		0	6 months 2°C 29°C	6 months 2°C	29°C
Nonwaxy					
Milled	dry	36.6	41.7	41.7	—
Milled	wet	26.5	31.9	34.1	39.6
Rough	wet	26.5	40.6	45.8	37.5
Defatted milled ^c	wet	20.6	20.3	21.6	49.6
Waxy^d					
Milled	dry	26.6	24.8	22.4	—
Milled	wet	26.7	26.7	25.9	—

* Percent of powder retained in 80-mesh sieve after grinding in a Wig-L-Bug amalgamator for 20 s; LSD (5%) ≤ 2.4%.

^b LSD (5%) ≤ 0.7%.

^c Values are mean of two rices.


^d One rice line at each season.

(From ref. 17)

表四、不同貯存期及貯存方式下白米透明度之平均值

Table 4. Means of translucency under different storage periods and storage forms (From ref. 2)

Variety	Storage period (months)	storage form (rice from 1st crop)			
		cold storage	sack storage	bulk storage (with fan)	
TNG-67	0	3.0	3.0	3.0	3.0
	3	3.0	3.0	3.5	3.5
	6	3.0	3.0	4.0	4.0
	9	3.0	3.0	4.0	4.0
	12	3.0	4.0	4.0	4.0
Sen-3	0	3.0	3.0	3.0	3.0
	3	3.0	3.0	3.0	3.0
	6	3.0	3.0	3.0	3.0
	9	3.0	3.0	4.0	4.0
	12	3.0	3.5	4.0	4.0



圖一 貯藏期間白米黃變米率之變化

Fig. 1. Change in the percentage of discolored grain as affected by different treatments during storage.
(From ref. 2)

關於碾米品質的改變，主要是完整米率 (head rice yield) 的增加，由於貯藏後澱粉顆粒的微束鍵結增強，加上蛋白質的 S-H 基氧化成 S-S 基，使得米粒的硬度指數 (hardness index) 變大，故在碾米過程中有較大的抗力，避免斷裂，因此而增加了完整米率⁽¹⁷⁾ (表 5)。

表五、不同貯藏條件下 (2°C 及 29°C) 貯藏 6 個月後之非糯性糙米的完整米率和硬度變化

Table 5. Changes in head rice yield and hardness index of nonwaxy brown rice stored at 2°C and at 29°C for 6 months.

Rice	Head rice yield (% of milled rice)		Hardness index*	
	2°C	29°C	2°C	29°C
IR 1529-680-3	15	64	36.5	42.2
IR 480-5-9	28	60	44.0	50.3
IR 24	79	72	41.4	44.9

* Percent of powder retained in 80 mesh sieve after grinding in a Wig-L-Bug amalgamator for 20 s. LSD (5%) = 1.5%. (From ref. 22)

三、貯藏對米粒烹調品質的影響

米粒貯藏後，由於脂肪的水解、氧化，其組成分含量發生變化，游離脂肪酸的含量，室溫貯藏較低溫貯藏有增加的現象⁽²¹⁾ (表 6)，其中又以白米方式貯藏增加最多⁽¹⁴⁾ (圖 2)，故浸過舊米的水其 pH 值會降低，且貯藏愈久 pH 值降低愈多，因此可利用這種方法來鑑定新舊米。


表六 貯藏期間白米之脂肪組成的變化

Table 6. Change in lipid compositions of polished rice during storage

	g/100g of rice*		g/100g of extraneous lipid			
	Extra-neous lipid	Fat-by-hydrolysis	Phospholipid	Free fatty acids	Neutral fat	Unsaponifiable matter
Koshiji-wase 9°C (Niigata)	0.35	0.53	1.5	30.6	49.9	5.7
Room temp.	0.34	0.54	0.4	51.9	30.8	6.0
Asahi (Okayama)	0.35	0.55	1.5	32.0	54.9	5.6
Room temp.	0.36	0.54	0.6	51.6	33.0	4.9


* Moisture content: 13%. (From ref. 22)

再則因游離脂肪酸的增加提高了連續粘度變化圖譜 (amylogram) 的最大粘度 (maximum viscosity) (圖 3、圖 4)。此外利用 X-ray 繞射 (diffraction) 後之 X-ray 種類的不同，亦證明了游離脂肪酸的增加是提高最大粘度的主要因素⁽²²⁾ (圖 5、6、7)。




圖二 貯藏期間脂肪酸度的變化

Fig. 2. Changes in fat acidity during storage. (Norin No. 29)
(From ref. 14)




圖三 不同貯藏溫度下所得稻米之米粉和去脂米粉之連續粘度變化圖譜

Fig. 3. Amylograms of rice powder and defatted rice powder prepared from rice grains having different storage temperatures. (From ref. 22)




圖四 添加稻米抽出液對去脂米粉之連續粘度變化圖譜之影響

Fig. 4 Effect of rice extract added back to defatted rice on amylogram. (From ref. 22)



圖五、31°C 貯藏下米粉之X線模式


Fig. 5. X-ray patterns of rice powders taken at 31°C.
..... raw starch) (From ref. 22)



圖六 可使粘度增加之溫度下貯藏之米粉的X線模式


Fig. 6 X-ray patterns of rice powders taken at temperature of viscosity increase. (From ref. 22)

但 Shin et al.⁽¹⁵⁾ 則認為米粒貯藏後，最大粘度的增加是由於澱粉酶 (amylase) 活性的降低所引起，並非由游離脂肪酸的增加所造成（圖8、表7）。



圖七 達最大粘度時之溫度下貯藏之米粉的X線模式

Fig. 7 X-ray patterns of rice powders taken at temperature of maximum viscosity. (From ref. 22)



圖八、35°C貯藏下各種糙米之連續粘度變化圖譜

Fig. 8 Amylograms of undefatted brown rice flour (UBR) (A). ether defatted brown rice flour (EDBR) (B). methanol extracted ether-defatted brown rice flour (MEDBR) (C). and brown rice starch (D). during storage of brown rice at 35°C.
 —, flours or starch at storage time of 0 month
, flours or starch at storage time of 12 months
 (From ref. 15)


表七、35°C貯藏下各種不同糙米米粉之澱粉酶活性

Table 7. Amylose activity in various brown rice flours during storage of brown rice at 35°C

Type of flour ^a	Amylose activity (mg maltose/min/ml extracts)	
	Before storage	After 12 months
Undefatted brown rice flour (UBR)	0.206±0.012	0.114±0.009
Ether defatted brown rice flour (EDBR)	0.170±0.015	0.090±0.009
Methanol extracted ether defatted brown rice flour (MEDBR)	Nil	Nil

^a Mean±S.D. based on 3 samples. (From ref. 15)

碱性擴散程度 (alkali spreading scores) 因貯藏而有降低的趨勢，此暗示着膠化溫度 (gelatinization temperature) 可能會因貯藏而升高⁽²⁾ (圖 9)。加熱吸水率及膨脹容積也因溫度及貯藏時間的增加而增加，可溶性固形物則有減少的趨勢⁽¹⁴⁾ (表 8)，所以煮飯的適當時機就加長了。Pushpamma and Reddy⁽¹⁰⁾ 指出貯存 6 個月的舊米之適當炊飯時間較新收穫的米多 4 ~ 6 分鐘。



圖九、貯藏期間白米碱性擴散程度之變化

Fig. 9 Change in alkali spreading scores of milled rice during storage.
(From ref. 2)

表八、貯藏稻米之炊飯品質


Table 8. Cooking quality of stored rice*

	Sample	Expanded volume (ml)	Water uptake ratio	pH of residual liquid
Manryo	Brown rice storage	39.6	3.08	6.72
	Milled rice storage	38.4	3.05	5.66
	Control (4°C)	36.2	2.64	6.98
Koshisakae	Brown rice storage	33.8	2.66	6.33
	Milled rice storage	35.2	2.77	5.54
	Control (4°C)	32.6	2.51	6.70
Norin No.29	Brown rice storage	33.6	2.85	6.75
	Milled rice storage	32.5	2.76	6.18
	Control (4°C)	32.2	2.54	6.85

* Milling yield, 90%, After 50 days storage. (From ref. 14)

四、貯藏對食用品質的影響

稻米的食用品質主要受米飯的質地及香味的影響。一般直鏈性澱粉含量較高之秈稻 (indica rice)，煮成米飯時較粗糙而不具粘度。直鏈性澱粉含量較低的梗稻 (japonica rice) 煮成米飯時乾濕度適中，粘性中等而較鬆軟，而糯稻 (waxy rice) 則幾乎不含直鏈性澱粉，煮成飯時較粘而濕潤^(10,11)。通常貯藏後的非糯性稻米 (non-waxy rice) 煮成飯時，其硬度(hardness) 會有增加的現象，膠體軟硬度 (gel consistency) 隨貯藏時間的增加而變硬，粘性 (stickiness) 也有降低的趨勢。但糯米因幾乎不含直鏈性澱粉，故貯藏後，不論是硬度、膠體軟硬度或粘性皆無明顯的變化^(10,17,2,16)。(圖10，表9、10、11)



圖十、貯藏期間膠體軟硬度之變化

Fig. 10 Change in gel consistency of milled rice during storage.(From ref. 2)

表九、貯藏期間不同品種白米之米飯性質的變化

Table 9. Mean changes in cooked rice properties during storage of milled rice of six nonwaxy varieties at 28-30°C^a

Storage Period (mo)	Cooked Rice Instron Hardness (kg)	Gel Consistency (mm)	Amylograph		Viscosity (BU)
			Peak	Final at 94°C	Cooled to 60°C
0	7.4	65	541	359	703
1	7.5	60	592	379	750
2	8.4	54	620	400	793
3	8.8	53	652	440	820
4	8.8	52	649	426	835
5	8.6	50	678	441	851
6	8.4	56

* Source: Perez and Juliano (1982): used by permission.

表十、不同貯藏條件下米飯之質地變化

Table 10. Texture of cooked rice prepared from 4°C-stored and 40°C stored rice samples

Property	4°C-stored rice	40°C-Stored rice	t-Test ^b
Texture of cooked rice ^a			
Hardness	2.92 ± 0.145	3.49 ± 0.162	*
Stickiness	0.51 ± 0.054	0.10 ± 0.027	**
Cohesiveness	0.681 ± 0.0440	0.715 ± 0.0341	

^a Texturometer unit.

^b *p=0.01 and **p=0.001. (From ref. 16)


表十一、不同貯藏條件下 (2°C和29°C) 貯藏 6 個月後之各種白米的粘度變化

Table 11. Stickiness of IR24, IR29, and IR833-6-2 milled rice after storage for 6 months at 2°C and at 29°C.

Variety or line	Form of stored rice	Crop season	Stickiness (g/10 grains)*	
			2°C	29°C
Nonwaxy				
IR24	Milled	dry	196	96
IR24	Milled	wet	173	83
	Defatted milled	wet	168	106
	Rough	wet	128	114
Waxy				
IR29	Milled	wet	208	262
IR833-6-2	Milled(7.4%protein)	dry	168	160
	Milled(9.3%protein)	dry	127	120

* LSD (5%)=18. (From ref. 17)

由貯藏過的米粒煮成的米飯其蒸氣經由氣相層析儀 (Gc gas chromatography) 分析的結果，發現約有一百多種揮發性物質⁽¹⁶⁾ (圖11)，而主要是揮發性羧基化合物。又貯藏在40°C的米粒其不飽和脂肪酸：亞麻油酸 (linoleic acid) 及亞麻脂酸 (linolenic acid) 有減少的趨勢⁽²⁰⁾ (表12)。相對地，羧基化合物中之己烷醛(hexanal)有顯著的增加⁽¹⁶⁾ (表13)，其峯面積 (peak area) 由 95.2 增加到 204.1。此意味着在貯藏期間不飽和脂肪酸自行氧化後增加了羧基化合物的含量。而且認定己烷醛是造成稻米腐敗氣味的主要因子。



圖十一 米飯之上部揮發性物質的氣相層析圖譜

Fig. 11 Gas chromatograms of headspace volatiles of cooked rice
 (A). 4°C-stored rice; (B). 40°C-stored rice. (From ref. 16)

表十三、米飯之主要揮發性組成分的相對含量

Table 13 Relative amounts of major volatile components of cooked rice

Peak no.	Compound	Relative amount of volatile component			
		Headspace analysis ^a		SDE method ^b	
		4°C-Stored	40°C-Stored	4°C-Stored	40°C-Stored
Alkanals					
15 Pentanal		13.0 (4.4)	27.0 (5.7)	4.2	12.4
24 Hexanal		95.2 (31.8)	204.1 (43.2)	65.9	226.6
35 Heptanal		5.8 (2.0)	7.9 (1.7)	7.4	16.6
50 Octanal		6.9 (2.3)	6.1 (1.3)	18.2	24.8
63 Nonanal		11.6 (3.9)	8.7 (1.8)	40.9	48.9
76 Decanal		6.6 (2.2)	3.4 (0.7)	14.4	19.7
Alkenals					
41 trans-2-Hexenal		0.6 (0.2)	1.3 (0.3)	1.3	3.6
55 trans-2-Heptenal		3.6 (1.2)	6.6 (1.4)	13.7	47.0
66 trans-2-Octenal		3.4 (1.1)	5.9 (1.2)	22.2	63.3
81 trans-2-Nonenal		1.0 (0.3)	2.0 (0.4)	12.2	38.1
106 trans-2-trans-4-Decadienal		1.9 (0.6)	1.2 (0.3)	132.5	193.4
Aromatic aldehydes					
79 Benzaidehyde		1.0 (0.3)	3.0 (0.6)	4.7	16.5
95 Phenylacetaldehyde		0.5 (0.2)	0.9 (0.2)	18.7	43.4
Ketones					
7 Acetone		12.0 (4.0)	15.3 (3.2)	—	—
34 2-Heptanone		2.3 (0.8)	6.1 (1.3)	1.7	10.8
49 2-Octanone		1.5 (0.5)	1.4 (0.3)	2.8	4.1
62 2-Nonanone		0.3 (0.1)	0.6 (0.1)	1.5	3.7
56 6-Methyl-5-hepten-2-one		1.4 (0.5)	2.3 (0.5)	2.0	11.9
Alcohols					
43 1-Pentanol		11.5 (3.8)	9.1 (1.9)	30.5	44.9
57 1-Hexanol		26.1 (8.7)	10.3 (2.2)	127.1	93.3
69 1-Heptanol		1.2 (0.4)	1.5 (0.3)	12.4	23.9
82 1-Octanol		1.5 (0.5)	1.8 (0.4)	23.0	41.4
96 1-Nonanol		0.6 (0.2)	0.3 (0.1)	12.1	14.8
68 1-Octen-3-ol		2.2 (0.7)	5.3 (1.1)	9.1	40.9
112 Benzyl alcohol		0.4 (0.1)	0.8 (0.2)	14.6	35.0
Miscellaneous					
42 2-Pentylfuran		13.0 (4.4)	18.1 (3.8)	11.5	30.3
155 4-Vinylphenol		— —	— —	66.3	126.7
Total peak area		299.0	472.4	—	—

^a Relative amount of each volatile component in the headspace vapor of cooked rice:(A). Peak area on the gas chromatogram;(B). peak area % (total peak area =100).

^b Relative amount of each volatile component in the extract obtained by simultaneous distillation-extraction of boiled rice (peak area of each volatile component×100/peak area of internal standard). (From ref. 16)

表十二、貯藏稻米之脂肪組成

Table 12. Lipid composition of stored rice (From ref. 20)

Storage temperatures	% Composition of total lipid	
	5°C	40°C
Fatty acids	Myristic acid	0.6
	Palmitic acid	17.0
	Stearic acid	0.6
	Oleic acid	30.4
	Linoleic acid	47.1
	Linolenic acid	4.3
Total lipid (mg% of rice)	390	310


遠藤等人⁽⁴⁾指出由米飯質地分析儀 (texturometer) 測出之硬度、粘性及附着性 (cohesiveness) 與官能檢定 (sensory test) 的綜合評價有密切的關係，其中硬度與綜合評價呈顯著負相關，而粘性與附着性則與之有顯著正相關。(表14)

表十四、米飯質地分析儀所測定的特性值和官能品評之間的相關係數

Table 14. Correlation coefficients between texturometer measurements and sensory score (From ref. 4)

	Hardness	Cohesiveness	Hardness/ cohesiveness	Stic- kiness	Hardness/ stickiness	Mean stickiness
Sensory score	-0.50*	0.82**	-0.77**	0.81**	-0.73**	0.81**

由以上的結果，貯藏後米飯質地變硬，粘性變差，加上腐敗的氣味，可知舊米的食用品質降低，在食用上較不為品評員所接受^(19, 6, 20)(圖12)，當然也較不受大眾歡迎。



圖十二、貯藏 3 個月後稻米之官能評價

Fig. 12 Organoleptic evaluation of rice stored for 3 months (From ref. 19)

五、貯藏對營養品質的影響

稻米的營養品質，主要決定於蛋白質的含量，尤其是離胺酸 (Lysine) 含量的多寡。米粒經貯藏後，游離胺基酸 (free amino acid) 中除胺基丙酸 (alanine) 外，其他的胺基酸如：離胺酸、甲硫胺酸 (methionine)、麸胺酸 (glutamic acid) 等皆有降低的現象，而且隨貯藏溫度及時間而異，溫度愈高、時間愈長則游離胺基酸含量降低愈多⁽⁵⁾ (表15)。但水溶性及鹽溶性蛋白質的含量則幾乎沒有太大的改變^(17, 9) (表16)。

表十五、貯藏期間米粒之游離胺基酸含量的變化

Table 15 Change in free amino acids of rice grains (Nippon bare) during storage under air (mg/100g dry wt.) (From ref. 15)

Amino acid	Before storage	1 year		3 year	
		Paddy 30°C	Pady 4°C	Paddy 4°C	Brown 4°C
Lysine	16.5	10.1	13.4	1.8	1.9
Histidine	4.8	3.2	4.7	0.7	0.9
Arginine	14.9	10.5	14.0	3.8	4.4
Aspartic acid	32.6	23.8	21.9	7.4	8.3
Threonine	10.5	5.1	7.2	2.0	2.0
Serine	15.9	10.2	12.3	7.1	9.2
Glutamic acid	58.3	36.8	45.4	7.2	10.3
Proline	10.6	7.2	8.1	1.9	1.9
Glycine	5.6	3.7	3.9	2.0	1.7
Alanine	10.0	12.2	10.9	14.4	10.8
Valine	8.2	4.8	5.7	2.5	2.4
Methionine	8.1	5.1	6.1	1.5	1.3
Isoleucine	4.6	2.7	3.0	0.7	0.6
Leucine	9.0	5.1	5.9	1.1	0.9
Tyrosine	5.7	3.3	3.7	1.0	0.8
Phenylalanine	4.5	2.6	3.0	0.7	0.7
Total	219.8	146.4	169.2	55.8	58.1

表十六 用 MAES 法所抽出 albumin 與 globulin 量

Table 16. The amount of albumin and globulin by maes method (From ref. 9)

Material	Albumin % of total N	Globulin % of total N
Check	3.80	6.92
Stored rice	3.35	6.68
Heated rice	3.17	6.56

至於維生素 B₁ (即 thiamine) 的含量則因貯藏漏失而含量降低⁽²¹⁾ (表17)，且其降低速率較核黃素 (riboflavin) 及菸鹼酸 (niacin) 快⁽¹⁰⁾。由於離胺酸及維生素 B₁ 含量的降低而降低了米粒的營養品質。除此之外，稻米在貯藏期間，由於微生物的滋長，產生黴菌毒素 (mycotoxins) (由真菌類 (fungi) 所產生)，其中之黃麴毒素 (aflatoxin) (由黴菌 (mold) 所產生) 對人類或是動物更是為害很大⁽⁷⁾。

表十七、貯藏溫度對於白米中維生素B₁含量的影響

Table 17. Effect of storage Temperature on thiamine content of polished rice.

Variety	treatment	(r-%) Thiamin content
Koshijiwase (Niigata)	Before storage	66.2
	9°C	58.3
	Room Temp.	33.2
Asahi (Okayama)	Before storage	98.6
	9°C	68.0
	Room Temp.	57.7

(From ref. 21)

六、結論

由以上各點的說明可知；因為貯藏的關係，在米粒外貌、烹調、食用及營養品質上都有不良的變化，同時在貯藏期間易受微生物代謝物的污染，國民食用後易有礙健康，故貯藏後稻米品質劣化是不爭的事實。要解決這個問題，收穫後的處理要特別小心，如乾燥機之選別、乾燥溫度之調整及乾燥後穀粒之含水量均要在規定之範圍內。本省農會及碾米廠之稻米貯藏，以袋裝及散裝兩種方式為主。根據宋及洪⁽²⁾之研究，散裝者之品質劣化程度較袋裝者大，尤以無通風設備之散裝稻穀為甚。稻穀本身為有生命的種子，故在貯藏中亦進行呼吸作用並消耗能源。在空氣不流通而高溫的地方，則呼吸率增加，ATP 的消耗增加，種子之老化也增快。因此，如何降低溫度，增加空氣的流通，以減緩種子之老化，乃為保持稻穀新鮮度之要訣。目前臺灣各碾米廠使用加入 CO₂ 於包裝袋內之作法，可以維持稻米新鮮度相當久的時間，故值得大力推廣。日本的穀倉大部份有冷藏設備，故可以保持良質米的時間相當長，但在本省大規模的稻穀冷藏倉庫不多。政府大力推廣良質米行銷之際，興建新米倉時，應把冷藏設備考慮在內，如此才能長期保鮮本省的良質米。

參考文獻

- 宋勳 1978 臺中地區農會貯藏稻谷品質探討之試驗 臺中區農業改良場研究彙報新 2 : 17—25。
- 宋勳、洪梅珠 1987 貯藏方式對稻谷倉儲期間品質之影響 臺中區農業改良場研究彙報 (14、15) : 15—26。
- 林子清、陳賢哲 1984 日本稻米品質改良的研究 食品工業 16 (3) : 19~27。
- 遠藤勳、柳瀬肇、石間紀男、竹生新治郎 1980 極少量炊飯方式による米飯のテクスチュロメーター測定，食総研報，37 : 1—8。
- Aibara S, I. A. Ismail, H. Yamashita. 1986 Changes in rice bran lipids and

- free amino acids during storage Agri. Biol. Chem. 50(3): 665~673.
6. Calderwood D. L., R. R. Cogburn, B. D. Weble and M. A. Marchetti. 1984
Aeration of rough rice in long-term storage transactions of the ASAE pp.
1579~1585.
 7. Coghurn R. R. 1985 Rough rice storage pp. 265-288 In: Rice Chemistry and
Technology 2nd ed. B. O. Juliano ed. Am. Assoc. Cereal Chem. St. Paul MN.
 8. Eggum. B. O. 1979 The nutritional value of rice in composition with other
cereals pp. 91-111. In: Pro. Workshop on chmical Aspects of Rice grain
Quality. Irt. Rice Res. Los. Banos, Laguna Philippines.
 9. Iwasaki. T., N. Shihuya and S. Chikuhu. 1972 Studies on Rice Protein. J.
Jpn. Soc. Food. Sci. Technol (In Japanese) 19(2):70~75.
 10. Juliano. B. O. 1985 Criteria and tests for rice grain qualities pp. 443-524 In:
Rice Chemistry and Technology 2nd ed. B. O. Juliano ed. Am. Assoc. Cereal
Chem. St. Paul. MN.
 11. Juliano. B. O. 1979 The Chemical basis of Rice grain quality. pp. 69-90. In:
Pro. Workshop on Chemical Aspects of Rice Grain Quality. Irt. Rice Res.
Los Banos, Laguna Philippines.
 12. Mallick A. K. and B. Nandi. 1982 Deterioration of stored rough rice 5.
grain in private storage. Seed Sci & Technol 10:527-533
 13. Sharp R. N. and L. K. Timme. 1986 Effects of storage time temperature
and packaging method on shelf life of brown rice. Cereal Chem 63(3): 247-
251.
 14. Shibuya. N., T. Iwasaki, H. Yanase and S. Chikubu. 1974 Studies on det-
erioration of rice during storage J. Jpn. Soc. Food Sci. Technol. 21(12):597-
603. (In Japanese)
 15. Shin. M. G., J. S. Rhee and T. W. Knon. 1985 Effects of Amylase activity
on changes in Amylogram characteristics during storage of Brown Rice
Agri. Biol. Chem. 49(8):2505-2508.
 16. Tsugita T. T. Ohta and H. Kata. 1983 Cooking flavor and texture of Rice
stored under different conditions. Agri. Biol. Chem. 47(3): 543-549.
 17. Villareal R. M., A. P. Resurreuion, L. B. Suzuki and B. O. Juliano. 1976
Changes in physicochemical properties of rice during storage Die starke 28:
88-94.
 18. Yanai S., T. Ishitani and Y. Kakiuchi. 1979 Influence of gaseous environm-
ent on hermetic storage of brown rice. Nippon Shokuhin Kogyo Gakkaishi
26(1): 225-231. (In Japanese)
 19. Yanai S., T. Ishitani and T. Kozo. 1979 Influence of gaseous environment
on the hermetic storage of Milled rice. Nippon Shokuhin Kogyo Gakkaishi
26(3): 145-150.
 20. Yasumatsu K., S. Moritaka, S. Wada. 1966. Stadies on Cereals. Agri. Biol.
Chem. 30(5): 478-486.

21. Yasumatsu K. and S. Moritaka. 1964. Fatty acid compositions of rice lipid and their changes during storage. *Agri. Biol. Chem.* 28(5):257-264.
22. Yasumatsu K., S. Moritaka and T. Kakinuma. 1964. Effect of the change during storage in lipid composition of rice on its Amylogram. *Agri. Biol. Chem.* 28(5): 265-272.

Changes in Rice Grain Quality During Storage

L. H. Wang and S. C. Hsieh

National Chung Hsing University and Taichung District Agricultural
Improvement Station

ABSTRACT

Physico-chemical properties of rice is known to change during the storage period. The mechanism of aging during storage and the effect of storage time on grain appearance, milling quality, cooking and eating qualities as well as nutritional value are reviewed and evaluated.

Free fatty acid is increased in parallel with the increase of pH value. Free fatty acid and amylose are then forming the fatty acid-amylose complex which contributed very much to aging of rice grains. Gelatinization temperature, Water absorption rate and expansion volume are increased with the prolongation of storage period. Harder gel consistency and reduced stickiness of rice grains are resulted due to storage. Free amino acid and vitamin B₁ contents are reduced with the prolongation of storage period. However, no change in the rate of water and alkali soluble proteins occurred during the storage period.