表四、花蓮地區水稻品種間白葉枯病不同罹病率之糙米外觀（1998-2000）

Table 4. Effects of bacterial leaf blight severity on grain appearance characteristics in Hualien area for 1998-2000.

Location	Cultivar	Disease severity*	Grain appearance (1st crop)				Grain appearance (2nd crop)			
			White center	White belly	White back	Trans-lucency	White center	White belly	White back	Trans-lucency
Fenglin	TK2	Light	0.3**	0	0	3.0	0.5	0.2	0	3.3
		Medium	0.3	0	0	3.0	0.2	0.5	0	3.3
		Heavy	0.4	0	0	3.0	0.7	0.3	0	3.3
	TK9	Light	0.3	0	0	3.0	0.6	0.2	0	3.0
		Medium	0.3	0	0	3.0	0.7	0.2	0	3.2
		Heavy	0.8	0	0	3.0	0.9	0.3	0	3.2
	TK16	Light	1.0	0	0	3.0	0.8	0.1	0	3.2
		Medium	1.2	0	0	3.0	1.3	0.2	0	3.2
		Heavy	1.0	0	0	3.0	1.4	0.5	0	3.5
Yuli	TK2	Light	0.8	0	0	3.0	0.6	0	0	3.2
		Medium	0.8	0.1	0	3.0	0.6	0.1	0	3.2
		Heavy	1.0	0	0	3.0	0.7	0.2	0	3.2
	TK9	Light	1.0	0	0	3.0	0.5	0.2	0	3.0
		Medium	0.7	0.2	0	3.0	0.6	0.2	0	3.0
		Heavy	0.7	0.2	0	3.0	0.6	0.2	0	3.0
	TK16	Light	1.3	0.2	0	3.0	0.7	0	0	3.0
		Medium	1.3	0.2	0	3.0	1.0	0.1	0	3.0
		Heavy	1.3	0.2	0	3.0	1.0	0.5	0	3.0
Fuli	TK9	Light	1.0	0.4	0	3.0	0.6	0	0	3.0
		Medium	0.9	0.5	0	3.0	1.1	0	0	3.0
		Heavy	0.9	0.6	0	3.3	0.9	0.2	0	3.0
	TK16	Light	1.4	0.1	0	3.0	1.8	0.1	0	3.3
		Medium	1.5	0.1	0	3.0	1.8	0.2	0	3.3
		Heavy	1.4	0.2	0	3.0	1.9	0.4	0	3.3
	KH139	Light	0.5	0.9	0	3.0	0.1	0.7	0	3.0
		Medium	0.2	1.0	0	3.0	0.1	0.7	0	3.0
		Heavy	0.5	1.1	0	3.0	0.1	1.0	0	3.0

* Disease severity : Light (the injury rate of disease area on the flag leaf : 0~1/3).

Medium (the injury rate of disease area on the flag leaf : 1/3~2/3).

Heavy (the injury rate of disease area on the flag leaf : over 2/3).

** Each value is the mean of 1998-2000.