表六、花蓮地區水稻品種間白葉枯病不同罹病率對米飯食味表現（1998-2000）

Table 6. Effects of bacterial leaf blight severity on panel taste value characteristics in Hualien area for 1998-2000.

Location	Cultivars	Diseaseseverity*	Panel taste value**
Fenglin	TK2	Light	3B1C
		Medium	3B1C
		Heavy	3B1C
	TK9	Light	4B1C
		Medium	4B1C
		Heavy	2B3C
	TK16	Light	1A4B
		Medium	5B
		Heavy	4B1C
Yuli	TK2	Light	3B2C
		Medium	3B2C
		Heavy	2B3C
	TK9	Light	3B2C
		Medium	1B4C
		Heavy	2B3C
	TK16	Light	2B3C
		Medium	2B3C
		Heavy	2B3C
Fuli	TK9	Light	1A2B1C
		Medium	1A2B1C
		Heavy	3B2C
	TK16	Light	1A4B
		Medium	4B1C
		Heavy	3B2C
	KH139	Light	1A4B
		Medium	5B
		Heavy	3B2C

* Disease severity : Light (the injury rate of disease area on the flag leaf :0~1/3).
Medium (the injury rate of disease area on the flag leaf :1/3~2/30.
Heavy (the injury rate of disease area on the flag leaf :over 2/3).

**Grade B is the same with CK sample, A is superior and C is inferior to B.