

Table 3. Cumulative number of deaths and mortality in *Mikania micrantha* of differing ground diameters after 3 monthly cuttings in the summer. The 1st cutting was conducted on June 22, 2000

Diameter class (mm)	Sampled plants	Cumulative death and mortality		
		1st cutting	2nd cutting	3rd cutting
1.0~2	4	3 (75%) ¹⁾	4 (100%)	4 (100%)
2.1~3	12	3 (25%)	12 (100%)	12 (100%)
3.1~4	14	5 (36%)	10 (71%)	13 (93%)
4.1~5	11	9 (82%)	11 (100%)	11 (100%)
5.1~6	4	2 (50%)	4 (100%)	4 (100%)
6.1~7	3	1 (33%)	1 (33%)	3 (100%)
7.1~8	2	2 (100%)	2 (100%)	2 (100%)
Total	50	25 (50%)	44 (88%)	49 (98%)

1) The numbers in parentheses indicate the cumulative mortality.

Table 4. Cumulative mortality of *Mikania micrantha* after various stem cuttings in different seasons

Season	Date of first cutting	Cumulative mortality (%)		
		1st cutting	2nd cutting	3rd cutting
Summer	June 22, 2000	50	88	98
Autumn	Sept. 28, 2000	50	92	92
Winter	Dec. 30, 2000	30	44	60
Spring	Mar 29, 2001	4	24	52