



圖七、巴克素處理後 *Phal. Sogo Kitty* 開花表現，CK:對照組、A:抽梗前  $0.25 \text{ mL}\cdot\text{L}^{-1}$  、B:抽梗前  $0.125 \text{ mL}\cdot\text{L}^{-1}$  、C:抽梗後  $0.5 \text{ mL}\cdot\text{L}^{-1}$  、D:抽梗後  $0.25 \text{ mL}\cdot\text{L}^{-1}$  及 E:抽梗後  $0.125 \text{ mL}\cdot\text{L}^{-1}$

Fig. 7. Flowering performance of *Phal. Sogo Kitty* after paclobutrazol treatments. CK:control, A:before  $0.25 \text{ mL}\cdot\text{L}^{-1}$ , B: before  $0.125 \text{ mL}\cdot\text{L}^{-1}$ , C: after  $0.5 \text{ mL}\cdot\text{L}^{-1}$ , D: after  $0.25 \text{ mL}\cdot\text{L}^{-1}$  and E: after  $0.125 \text{ mL}\cdot\text{L}^{-1}$ .



圖八、三種蝴蝶蘭品種經不同巴克素處理後開花表現，A: *Phal. Arco Little Caroline*、B: *Phal. Jiuhbao Pink Lady* 及 C: *Phal. Ney Shan Gu Niang*

Fig. 8. Flowering performance of paclobutrazol treatments. A: *Phal. Arco Little Caroline*, B: *Phal. Jiuhbao Pink Lady* and C: *Phal. Ney Shan Gu Niang*.