

表七、102 年一期作礁溪鄉及 102 年二期作竹田村水稻田雜草受施用米糠之影響

Table 7. Weights of weeds treatment with rice bran at the first crop in Jiaosi and second crop in Chuten area in 2012.

Species of weeds	Treatment ^z				C.K.			
	N	Fresh weight (g)	Dry weight (g)	Relative weight (g) ^y	N	Fresh weight (g)	Dry weight (g)	Relative weight (g)
Jiaosi(102-I)								
<i>Monochoria vaginalis</i> (Burm. F.) Presl	-	-	-	-	-	-	-	-
<i>Lindernia procumbens</i> (Krock.) Borbas	85	70.89	8.28	0.83	163	188.48	32.28	1.16
<i>Echinochloa crusgalli</i> (L.) P. Beauv.	-	-	-	-	-	-	-	-
<i>Sphenoclea zeylanica</i> Gaertn.	-	-	-	-	-	-	-	-
<i>Paspalum distichum</i> L.	-	-	-	-	-	-	-	-
<i>Fimbristylis miliacea</i> (L.) Vohl	-	-	-	-	-	-	-	-
<i>Cyperus iria</i> L.	-	-	-	-	-	-	-	-
<i>Ludwigia hyssopifolia</i> (G. Don) Exell.	-	-	-	-	-	-	-	-
Chuten(102-II)								
<i>Monochoria vaginalis</i> (Burm. F.) Presl	-	-	-	-	1	1.37	0.07	1.37
<i>Lindernia procumbens</i> (Krock.) Borbas	7	5.80	0.63	0.83	33	49.89	4.93	1.51
<i>Echinochloa crusgalli</i> (L.) P. Beauv.	-	-	-	-	4	2.62	0.44	0.66
<i>Sphenoclea zeylanica</i> Gaertn.	34	101.62	8.50	2.99	76	165.62	13.83	2.18
<i>Paspalum distichum</i> L.	-	-	-	-	3	1.48	0.25	0.49
<i>Fimbristylis miliacea</i> (L.) Vohl	1	1.26	0.14	1.26	38	77.12	8.47	2.03
<i>Cyperus iria</i> L.	-	-	-	-	13	76.47	8.07	5.88
<i>Ludwigia hyssopifolia</i> (G. Don) Exell.	-	-	-	-	2	3.87	0.46	1.93

^zMeans of 1,000 kg/ha of rice bran application amount.^yMeans of 100%* fresh weight/weeds number(N).